

ORİJİNAL ARAŞTIRMA ORIGINAL RESEARCH

DOI: 10.31609/jpmrs.2019-72299

The Role of Attention- Deficit/Hyperactivity Disorder in the Etiology of Osgood Schlatter Disease

Osgood Schlatter Hastalığının Etyolojisinde Dikkat Eksikliği/Hiperaktivite Bozukluğunun Rolü

 Banu ORDAHAN^a,  Hatice UĞURLU^a

^aNecmettin Erbakan University Meram Faculty of Medicine, Department of Physical Medicine and Rehabilitation, Konya, TURKEY

ABSTRACT Objective: Osgood-Schlatter disease (OSD) is an overuse injury and regular practice of sports activity is a major risk factor for OSD. However, OSD is not uncommon in children who do not participate in sports in routine orthopedic practice. Over activity is not always related with sports but may be related with Attention-Deficit/Hyperactivity Disorder (ADHD). This study aimed to investigate the symptoms of ADHD in children with OSD. **Material and Methods:** Forty five male patients, with Osgood-Schlatter disease, and 45 age-matched healthy volunteers were included in the study. Symptoms of ADHD of the volunteers were evaluated using Conners' Parent Rating Scale (CPRS-R:S). Both groups' functional status was evaluated using Pediatric Quality of Life Inventory (PedsQL). **Results:** In the patient group, a statistically significant increase was found in Hyperactivity and Oppositional subscales of Conners' Parent Rating Scale- Revised Short compared with the control group ($p < 0.05$). Pediatric Quality of Life Scale was significantly lower in the patient group compared with the control group ($p < 0.05$). **Conclusion:** Our findings suggest that ADHD is a significant risk factor for OSD. Especially during the evaluation of OSD patients, who do not engage in sports activity, physicians should be aware of ADHD symptoms and refer children to psychiatrists for additional evaluations.

Keywords: Osgood-Schlatter disease;
Attention-Deficit/Hyperactivity Disorder;
hyperactivity and impulsivity

ÖZET Amaç: Osgood-Schlatter hastalığı (OSH), aşırı kullanım yaralanmasıdır ve düzenli spor aktiviteleri OSH için önemli bir risk faktörüdür. Bununla birlikte, rutin ortopedik pratikte spora katılmayan çocuklarda OSH nadir değildir. Aşırı aktivite her zaman sporla ilgili değildir, ancak Dikkat Eksikliği/Hiperaktivite Bozukluğu (DEHB) ile ilişkili olabilir. Bu çalışmada, OSH olan çocuklarda DEHB semptomlarını araştırmayı amaçladık. **Gereç ve Yöntemler:** Çalışmaya OSH'li 45 erkek hasta ve yaş uyumlu 45 sağlıklı gönüllü alındı. Gönüllülerin DEHB belirtileri, Conners'in Ebeveyn Değerlendirme Ölçeği (CPRS-R:S) kullanılarak değerlendirildi. Her iki grubun fonksiyonel durumu Pediatrik Yaşam Kalitesi Envanteri (PedsQL) kullanılarak değerlendirildi. **Bulgular:** Hasta grubunda, kontrol grubuna kıyasla Conners Ebeveyn Değerlendirme Ölçeği, hiperaktivite ve karşı koyma alt ölçeklerinde istatistiksel olarak anlamlı bir artış bulundu ($p < 0,05$). Pediatrik Yaşam Kalitesi Ölçeği skorları hasta grubunda kontrol grubuna göre anlamlı derecede düşüktü ($p < 0,05$). **Sonuç:** Bulgularımız DEHB'nin OSH için önemli bir risk faktörü olduğunu göstermektedir. Özellikle spor aktivitesiyle uğraşmayan OSH'li hastaların değerlendirilmesi sırasında, doktorlar DEHB belirtilerinin farkında olmalı ve ek değerlendirmeler için çocukları psikiyatristlere yönlendirmelidir.

Anahtar Kelimeler: Osgood-Schlatter hastalığı;
Dikkat Eksikliği/Hiperaktivite Bozukluğu;
hiperaktivite ve dürtüsellik

Osgood-Schlatter's disease (OSD) is an overuse injury caused by tension on the patellar tendon placed on the anterior tibial tubercle. It presents with local pain, swelling and tenderness over the tibial tuberosity and most frequently occurs in girls between the age of 11 to 13 years and in boys between 13 and 15 years. The symptoms are usually exacerbated with

sporting activities that especially involve jumping, running or stair climbing.¹⁻³ Radiologic studies have led to the theory that OSD may be caused by chronic tensile stress on the patellar tendon within the growth plate, which leads to irritation and inflammation. Inflammation results from micro-trauma to the growth plate with repetitive use before closure of the epiph-

Correspondence: Banu ORDAHAN
Necmettin Erbakan University Meram Faculty of Medicine,
Department of Physical Medicine and Rehabilitation, Konya, TURKEY/TÜRKİYE
E-mail: banuordahan@gmail.com

Peer review under responsibility of Journal of Physical Medicine and Rehabilitation Science.

Received: 08 Nov 2019

Received in revised form:

Accepted: 13 Dec 2019

Available online: 10 Jan 2020

1307-7384 / Copyright © 2020 Turkey Association of Physical Medicine and Rehabilitation Specialist Physicians. Production and hosting by Türkiye Klinikleri.

ysis, causing traction injury and microavulsions of the tubercle.³⁻⁵

Attention-Deficit/Hyperactivity Disorder (ADHD) is a heterogeneous neurodevelopmental disorder characterized by the co-existence of inattention, impaired cognitive function, lack of coordination, hyperactivity, and impulsivity which cause functional impairment. It is one of the common psychiatric disorders among children with an estimated prevalence between 6.7 and 7.8% of all children in school age.^{6,7}

We hypothesized that ADHD might be another risk factor for those who do not participate in sports. The purpose of this study was to investigate the relationship between OSD and ADHD.

MATERIAL AND METHODS

Forty five male patients with OSD, aged between 9-15 years, and 45, age- matched healthy controls were included in the study. The study was approved by the local research ethics committee of Necmettin Erbakan University Medical Faculty. The diagnosis of OSD was made based on patient history and physical examination findings. To confirm the diagnosis of OSD, the following diagnostic criteria were applied:^{3,8} (1) pain localized to the tibial tubercle, especially after severe physical activity, (2) tenderness, erythema and local swelling and over the tibial tubercle, (3) pain should be present on the tibial tubercle resulting from forced knee extension. Knee radiographs were taken in all patients. Radiographs may illustrate OSD and are confirmatory for the diagnosis. Lateral radiographs can show prominence of the tibial tuberosity with anterior ossification separating from the proximal tibia.^{3,8} The presence of ADHD was assessed using Conners' Abbreviated Rating Scale (CARS) answered by the parents.

Conners Parent Rating Scale-Revised Short (CPRS-R:S):CPRS-R:S was created by Kumar et al. The CPRS-R:S is a 27-item questionnaire divided into four subscales: opposition, cognitive problems/inattention, hyperactivity, and ADHD index.^{9,10} Each item is rated on a four-point scale, from 0 (not true at all) to 3 (very much true). The Turkish version of the CPRS was studied by Kaner et al.¹⁰

Pediatric Quality of Life Inventory (PedsQL): which has been used in this study has 23 questions including physical functioning, emotional functioning, social and school functioning, to assess health-related quality of life in children.¹¹ Five choice Likert type visual scale is used in the inventory (0=never, 1=almost never, 2=sometimes, 3=often, 4=almost always). Points obtained from the items are converted linearly to a value ranging between 0 and 100 (0=100, 1=75, 2=50, 3=25, 4=0). Higher scores reflect higher quality of life. The reliability and validity of the Turkish PedsQL 8-12 and 13-18 age group forms have been studied by Memik et al. and reported to have high reliability and validity.^{12,13}

STATISTICAL ANALYSIS

The SPSS for Windows 15.0 software package . (SPSS for Windows, Version 16.0. released 2007; SPSS Inc.,USA) was used for the statistical evaluation of the data. Conformity of continuous variables with normal distribution was investigated using the Kolmogorov-Smirnov test. All variables were distributed normally. Descriptive data are presented as mean \pm standard deviation. Demographic and clinical characteristics were compared using the Chi-square test. The independent samples t-test was used to compare the two groups. The significance level was accepted as $p < 0.05$.

RESULTS

The baseline characteristics of the patients are given in Table 1. In the patient group, a statistically significant increase was found in Hyperactivity and Opposition subscales of CPRS-RS compared with the control group ($p < 0.05$) (Table 2). The PedsQLS was significantly lower in the patient group compared with the control group ($p < 0.05$) (Table 2).

TABLE 1: Demographic characteristics of the patients.

	Study group (n=45)	Control group (n=45)	p
Age (years)	12.20 \pm 3.08	11.80 \pm 3.24	0.82
BMI (kg/m ²)	20.01 \pm 2.36	21.12 \pm 2.98	0.86

Data are given as mean \pm SD, BMI: Body Mass Index.

TABLE 2: Assessment of Conners' Parent Rating Scale-Revised Short and Pediatric Quality of Life Scale.

CPRS-RS	Study group	Control group	P value
Oppositional subscale	7.01±4.45	5.22±4.28	0.032
Cognitive problem subscale	5.14±4.21	4.78±3.98	0.720
Hyperactivity subscale	9.80±2.65	5.29±2.89	0.016
ADHD Index	16.02±5.05	13.82±4.95	0.021
PedsQL	42±10.02	82.2±11.32	0.001

Data are given as mean±SD, CPRS-RS: Conners' Parent Rating Scale-Revised Short; PedsQL: Pediatric Quality of Life Scale.

DISCUSSION

In this study, the relationship between ADHD and OSD was investigated. Several studies have examined the risk factors for Osgood-Schlatter disease. Lucena et al. showed that OSD was caused by shortening of the rectus femoris muscle.¹⁴ Shortening of the rectus femoris may substantially affect the biomechanical function of the knee. Demirbag et al. evaluated patellar tendon insertions in 20 patients with OSD and 15 healthy children using knee magnetic resonance imaging (MRI). Their study showed that if the patellar tendon attachment was more proximal and in a broader area to the tibia, this might play a role as a predisposing mechanical factor in the onset of OSD.¹⁵ Gigante et al. studied the relationship between Osgood-Schlatter disease and torsional abnormalities of the lower limb. They evaluated 21 patients with OSD and 20 healthy controls using computed tomography (CT). They found that increased external tibial torsion might be a risk factor in OSD.¹⁶ Nakasew reported that increased quadriceps femoris muscle tightness and muscle strength during knee extension and flexibility of the hamstring muscles were risk factors for OSD.¹⁷ Sarcević evaluated the relationship between limited dorsiflexion of the ankle and the occurrence of OSD in sports-active children. The author reported that limited dorsiflexion of the ankle joint might be of significant importance for OSD. The limitation of dorsiflexion within the ankle joint is associated with a compensatory increased knee flexion, tibial inversion, and foot pronation during the stance phase of running, thus increasing the stress on the patellar tendon.¹⁸

Most of these studies evaluated anatomic and mechanistic causes to clarify the pathophysiology of OSD. Current evidence indicates that OSD is an

overuse injury and regular practice of sports activity is major risk factor for OSD.¹ However, OSD is not uncommon in children who do not participate in sports in routine orthopedic practice. Over activity is not always related with sports but may be related with ADHD. Some authors reported that ADHD was a risk factor for acute traumatic injuries in children.¹⁹⁻²¹ Uslu et al. reported that children treated for extremity fractures would present with higher levels of impulsive-hyperactivity behavior problems than children treated for conditions other than fractures.¹⁹ Rowe revealed that children with ADHD were prone to all kinds of injury, more than the healthy population.²⁰ In a similar study, İz et al. showed that ADHD symptomatology was observed in most children admitted to the emergency department for accidental injuries.²²

Children with impulsive-hyperactivity behavior have been reported to have 1.5-2 times higher risk for injury and extremity fractures.²¹⁻²⁵

In a retrospective cohort study, Guy et al. assessed musculoskeletal injuries in patients aged under 17 years with ADHD. In the 11-year study period, the following traumatic musculoskeletal injuries were diagnosed in patients with ADHD; sprains (27.2%), fractures (21.2%), concussions (4.5%), dislocations (2.8%), and thoracic and pelvic injuries (0.6%). In this study, fractures were most frequently detected in the upper extremity injuries (60.5%), whereas sprains were most frequently detected in the ankle and foot injuries (36.0%). Among the categories of overuse injuries, specific arthropathies and connective tissue disorders have been identified in 29.1% of individuals with ADHD.²⁶

These studies showed that ADHD was a significant risk factor for acute injuries. However, we be-

lieve that children with ADHD are prone to both acute injuries and chronic repetitive injuries. Güler et al. investigated the relationship of OSD and ADHD. They found that ADHD was a significant risk factor for OSD; however no control groups were used in that study. The author recommended future case-control studies.⁸ In our study, there was a control group and the evaluation was performed in the male sex only. Therefore, we cannot generalize our results for both sexes. This is the most important limitation of our study. The assessment of ADHD symptoms was based on parent reports rather than psychiatric interviews. However, our main goal was to investigate the symptoms of ADHD in children with OSD. A statistically significant increase was found in favor of ADHD in hyperactivity and oppositional subscale

scores compared with the control group. There was also a statistically significant decrease in quality of life in the patient group compared with the control group.

CONCLUSION

During the evaluation of patients with OSD, physicians should be aware of ADHD symptoms and refer children to psychiatrists for additional evaluations.

Informed Consent

All procedures performed in studies involving human participants were obtained from all participants who participated in the study in accordance with the ethical standards of the institutional and national research committee and the 1964 Helsinki Declaration.

REFERENCES

1. Launay F. Sports-related overuse injuries in children. *Orthop Traumatol Surg Res* Feb. 2015;101(1 Suppl):139-47. PMID: 25555804
2. Indiran V, Jagannathan D. Osgood-Schlatter Disease. *N Engl J Med*. 2018;378:e15. PMID: 29539285
3. Whitmore A. Osgood-Schlatter disease. *JAAPA*. 2013;26:51-2. PMID: 24201924
4. Hirano A, Fukubayashi T, Ishii T, et al. Magnetic resonance imaging of Osgood-Schlatter disease: the course of the disease. *Skeletal Radiol*. 2002;31:334-42. PMID: 12073117
5. Gholve PA, Scher DM, Khakharia S, et al. Osgood Schlatter syndrome. *Curr Opin Pediatr*. 2007;19:44-50. PMID: 17224661
6. Thomas R, Sanders S, Doust J, et al. Prevalence of attention-deficit/ hyperactivity disorder: a systematic review and meta-analysis. *Pediatrics*. 2015;135:994-1001. PMID: 25733754
7. Leffa DT, Torres ILS, Rohde LA. A review on the role of inflammation in attention-deficit/hyperactivity disorder. *Neuroimmunomodulation*. 2018;25:328-33. PMID: 29874674
8. Guler F, Kose O, Koparan C, et al. Is there a relationship between attention- deficit/hyperactivity disorder and Osgood-Schlatter disease? *Arch Orthop Trauma Surg*. 2013;133:1303-7. PMID: 23748799
9. Kumar G, Steer RA. Factorial validity of the Conner's Parent Rating Scale-revised: short form with psychiatric outpatients. *J Pers Assess*. 2003;80:252-59. PMID: 12763699
10. Kaner S, Büyükoztürk S, İşeri E. Connors Anababa Dereceleme Ölçeği-yenilenmiş kısa: Türkiye standardizasyon çalışması. *Archives of Neuropsychiatry*. 2013;50:100-9.
11. Varn JW, Seid M, Rode CA. The PedsQL: measurement model for the pediatric quality of life inventory. *Med Care*. 1999;37:126-39. PMID: 10024117
12. Cakin Memik N, Ağaoğlu B, Coşkun A, et al. [The validity and reliability of the Turkish Pediatric Quality of Life Inventory for children 13-18 years old]. *Türk Psikiyatri Derg*. 2007;18:353-63. PMID: 18066726
13. Çakin Memik N, Ağaoğlu B, Coşkun A ve ark. Çocuklar için yaşam kalitesi ölçeğinin 8-12 yaş çocuk formunun geçerlik ve güvenirliği. *Türk J Child Adolesc Ment Health*. 2008;15:87-98.
14. de Lucena GL, dos Santos Gomes C, Guerra RO. Prevalence and associated factors of Osgood-Schlatter syndrome in a population-based sample of Brazilian adolescents. *Am J Sports Med*. 2011;39:415-20. PMID: 21076014
15. Demirag B, Ozturk C, Yazici Z, et al. The pathophysiology of Osgood-Schlatter disease: a magnetic resonance investigation. *J Pediatr Orthop B*. 2004;13:379-82. PMID: 15599229
16. Gigante A, Bevilacqua C, Bonetti MG, et al. Increased external tibial torsion in Osgood-Schlatter disease. *Acta Orthop Scand*. 2003;74:431-6. PMID: 14521294
17. Nakase J, Goshima K, Numata H, et al. Precise risk factors for Osgood-Schlatter disease. *Arch Orthop Trauma Surg*. 2015;135:1277-81. PMID: 26133498
18. Sarcević Z. Limited ankle dorsiflexion: a predisposing factor to Morbus Osgood Schlatter? *Knee Surg Sports Traumatol Arthrosc*. 2008;16:726-8. PMID: 18483807
19. Uslu M, Uslu R, Eksioğlu F, et al. Children with fractures show higher levels of impulsive-hyperactive behavior. *Clin Orthop Relat Res*. 2007;460:192-5. PMID: 17353797
20. Rowe R, Maughan B, Goodman R. Childhood psychiatric disorder and unintentional injury: findings from a national cohort study. *J Pediatr Psychol*. 2004;29:119-30.
21. Brehaut JC, Miller A, Raina P, et al. Childhood behavior disorders and injuries among and youth: a population based study. *Pediatrics*. 2003;111:262-9. PMID: 12563049
22. İz M, Çeri V. Prevalence of attention deficit hyperactivity disorder symptoms in children who were treated at emergency service due to unintentional injury. *Emerg Med Int*. 2018;2018:7814910. PMID: 30671264
23. Erdogan M, Desteli EE, Imren Y, et al. Is attention deficit and hyperactivity disorder a risk factor for sustaining fractures of proximal humerus? *Acta Chir Orthop Traumatol Cech*. 2014;81:221-6. PMID: 24945391
24. Kömürçü E, Bilgiç A, Hergüner S. Relationship between extremity fractures and attention-deficit/hyperactivity disorder symptomatology in adults. *Int J Psychiatry Med*. 2014;7:55-63. PMID: 24956917
25. Amiri S, Sadeghi-Bazargani H, Nazari S, et al. Attention-deficit/hyperactivity disorder and risk injuries: a systematic review and meta-analysis. *J Inj Violence Res*. 2017;9:95-105. PMID: 28554188
26. Guy JA, Knight LM, Wang Y, et al. Factors associated with musculoskeletal injuries in children and adolescents with attention-deficit/hyperactivity disorder. *Prim Care Companion CNS Disord*. 2016;18. PMID: 27733957.