

# Sonographic Evidence of Cheiralgia Paresthetica

## Cheiralgia Parestetikanın Sonografik Kanıtı

 Berke ARAS<sup>a</sup>,  Ömer KUZU<sup>a</sup>,  Emre ADIGÜZEL<sup>a</sup>,  Evren YAŞAR<sup>a</sup>

<sup>a</sup>Clinic of Physical Medicine and Rehabilitation, Ankara City Hospital, Ankara, Türkiye

**ABSTRACT** Cheiralgia paresthetica is the entrapment of the superficial branch of the radial nerve in the forearm. Although this nerve damage frequently occurs after direct trauma or compression, it may also be the result of repetitive forearm rotation movements. It occurs with symptoms such as pain, burning and tingling sensations in the distal forearm and hand. Electromyography can be used for diagnosis. Ultrasound is a feasible method of showing nerve compression. Although treatment is mostly conservative, surgical exploration is an option for resistant cases. The case is here presented of a 46-year old female agricultural worker with compression of the superior branch of the radial nerve due to occupation-related repetitive forearm rotation.

**ÖZET** Cheiralgia parestetika, radyal sinirin yüzeysel dalının ön kolda sıkışmasıdır. Bu sinir hasarı sıklıkla doğrudan travma veya kompresyon sonrası ortaya çıksa da tekrarlayan ön kol rotasyon hareketlerinin sonucu da olabilir. Hastalık distal ön kol ve elde ağrı, yanma ve karıncalanma hissi gibi bulgular ile ortaya çıkar. Tanıda elektromiyografiden yararlanılabilir. Ultrason, sinir sıkışmasını göstermenin uygun bir yöntemidir. Tedavi çoğunlukla konservatif olmakla birlikte dirençli olgularda cerrahi eksplorasyon bir seçenektir. Burada, meslekle ilişkili tekrarlayan ön kol rotasyonu nedeniyle radyal sinirin üst dalına bası olan 46 yaşında bir kadın tarım işçisi olgusu sunulmaktadır.

**Keywords:** Cheiralgia paresthetica; radial nerve; forearm pain

**Anahtar Kelimeler:** Cheiralgia parestetika; radyal sinir; ön kol ağrısı

Cheiralgia paresthetica, also known as Wartenberg syndrome, is a compression of the superficial radial nerve (SRN) in the distal forearm.<sup>1</sup> The SRN is a purely sensory nerve with no motor component. It supplies the skin on the lateral two-thirds of the dorsum of the hand and proximal parts of the lateral three and a half digits.<sup>2</sup> SRN entrapment may be caused by compression or direct trauma at any point along the nerve course. It is often compressed by tight bands (bracelet, watchstrap etc.) or a mass (lipoma, ganglion cyst, etc.).<sup>3</sup> In addition, with excessive, repetitive and long-lasting rotation movements of the forearm, the nerve can become compressed between the brachioradialis and extensor carpi radialis longus muscles.

The annual incidence of radial nerve compression syndromes is 0.003%.<sup>4</sup> The incidence of cheiralgia paresthetica, which is a subgroup of radial nerve entrapments, is unknown.<sup>5</sup> Cheiralgia paresthetica presents with complaints of pain, burning and numbness in the dorsolateral part of the distal forearm and hand. Often it is aggravated by activities such as hand rotation, pinching, and gripping. In addition to sensory symptoms, if there are findings such as muscle weakness, alternative diagnoses where the lesion may originate more proximal should be considered. It can be confused with deQuervain tenosynovitis due to its localization in the differential diagnosis of the disease. Ultrasound is an important method in demonstrating nerve compression in this disease. The

**Correspondence:** Berke ARAS

Clinic of Physical Medicine and Rehabilitation, Ankara City Hospital, Ankara, Türkiye

**E-mail:** drberkearas@gmail.com



Peer review under responsibility of Journal of Physical Medicine and Rehabilitation Science.

**Received:** 16 Jan 2022

**Received in revised form:** 30 Mar 2022

**Accepted:** 03 Apr 2022

**Available online:** 22 Apr 2022

1307-7384 / Copyright © 2022 Turkey Association of Physical Medicine and Rehabilitation Specialist Physicians. Production and hosting by Türkiye Klinikleri.

This is an open access article under the CC BY-NC-ND license (<https://creativecommons.org/licenses/by-nc-nd/4.0/>).

treatment of cheiralgia paresthetica is usually conservative (rest, splinting, nonsteroidal anti-inflammatory drugs and corticosteroid injection), but surgical exploration is an option for resistant cases.<sup>6,7</sup>

An uncommon case is here reported of SRN compression due to repetitive forearm rotation in a 46-year old female agricultural worker. It was diagnosed on the basis of clinical symptoms, physical examination, ultrasound and electromyography (EMG). To the best of our knowledge, SRN compression due to harvesting garlic has not been previously reported in the literature.

### CASE REPORT

A 46-year old female presented with the complaint of right-side anterior forearm pain, which had begun 3 days previously when she started to work harvesting garlic. It was learned that the work involved intense supination-pronation movements of the arm. The patient described pain, burning and numbness over the dorsolateral aspect of the distal forearm and hand in the distribution of the SRN, and she had difficulty in forearm rotation and gripping movements. There was no history of trauma or systemic disease. The physical examination revealed mild tenderness with palpation on the dorsolateral distal forearm and the Tinel test was positive (Figure 1). The ranges of motion of the elbow, wrist and metacarpophalangeal joints were full. There was no motor weakness or signs of atrophy. EMG and nerve conduction studies were consistent with sensory radial neuropathy with




FIGURE 1: The route of the superficial radial nerve at the wrist.

TABLE 1: Superficial radial nerve sensory conduction study parameters.

	Right-side	Left-side
SNAP CV (m/s)	41.6	51.5
SNAP amplitude (µV)	12.2	26.3

SNAP: Sensory nerve action potential; CV: Conduction velocity; m/s: Meter per second; µV: Microvolt.

no evidence of cervical radiculopathy (Table 1). In ultrasound imaging, in the area where the Tinel test and sonopalpation were most prominent, the diameter of the SRN was found to be larger than the same region of the other forearm (6 cm proximal to the radius styloid) (Figure 2). The patient was advised to rest and use a non-steroidal anti-inflammatory drug (acemetazine 60 m, twice daily). At the follow-up examination after 10 days, the patient’s complaints and examination findings had decreased significantly. Informed consent was obtained from the patient.




FIGURE 2: Sonographic evaluation of the superior branch of the radial nerve of the right (a) and left (b) forearm.

## DISCUSSION

In this case, SRN entrapment was seen to have developed in an agricultural worker engaged in harvesting garlic. It is thought that the SRN was compressed with the repeated supination-pronation movements required by the work. The increase in diameter in the nerve was revealed in the ultrasonographic examination, when compared with the contralateral arm.

After branching from the radial nerve in the antebrachial fossa, the SRN descends below the brachioradialis muscle and lateral to the radial artery. In the distal forearm, it pierces the deep fascia, and travels between the brachioradialis and extensor carpi radialis longus tendons. It then ramifies into dorsomedial and dorsolateral branches. These branches travel along with the abductor pollicis longus and the extensor pollicis brevis tendons in the first compartment of the wrist. While the dorsolateral branch supplies the radial aspect of the thumb, the dorsomedial branch innervates the region radial to the ulnar side of the ring finger.<sup>8</sup>

The SRN is sensitive to local trauma due to its superficial course.<sup>9</sup> In addition to tightly worn watches, bracelets or handcuffs, neuropathy may also develop with penetrating injuries. The SRN can also be damaged by external causes such as distal radius fracture and soft tissue masses (lipoma, ganglion cyst, etc.).<sup>10</sup> The SRN is frequently trapped in the area between the brachioradialis and extensor carpi radialis longus tendons. Especially with pronation movement, the extensor carpi radialis longus passes under the brachioradialis and compresses the nerve. Repetitive pronation and supination increase this compression and cause nerve damage.<sup>7</sup> In the current patient, the nerve was compressed due to repetitive pronation and supination required to harvest the garlic from the soil.

The severity of the disease varies according to the severity of the compression force and the duration of the compression. In mild cases, there is mild

pain and paraesthesia due to congestion and edema of the nerve, whereas in severe cases, irreversible allodynia and anesthesia can occur with arterial ischemia.<sup>11</sup> The nerve compression of the current patient was a reversible condition which responded well to rest and anti-inflammatory medication.

Diagnostic ultrasound is frequently used by physicians because of many advantages such as being easily accessible/applicable, non-invasive, economical, and repeatable, it does not involve ionizing radiation and has the sono-palpation feature. The SRN appears as parallel hyperechoic lines and hypoechoic structures between them are other peripheral nerves.<sup>12</sup> In previous studies, it has been reported that the cross-sectional area of the nerve is increased in compression neuropathies. It also gives information about other diseases to be considered in the differential diagnosis of wrist pain such as de Quervain tenosynovitis, scaphoid fracture and thumb carpometacarpal joint arthritis. In this case, no additional pathology other than cheiralgia paresthetica was found in the ultrasonographic evaluation. EMG is often performed for diagnosis, but it can be false negative and unable to show the exact site of the lesion.<sup>13</sup> In the current case, the patient's EMG findings were also consistent with the diagnosis of cheiralgia paresthetica.

In conclusion, cheiralgia paresthetica is a disease that should be kept in mind in patients with forearm-hand pain and paraesthesia. It can occur as a result of repetitive and long-lasting forearm rotations as in the current case. Measurement of the SRN diameter on ultrasound has very important diagnostic value in this disease.

### *Source of Finance*

*During this study, no financial or spiritual support was received neither from any pharmaceutical company that has a direct connection with the research subject, nor from a company that provides or produces medical instruments and materials which may negatively affect the evaluation process of this study.*

## REFERENCES

1. Tosun N, Tuncay I, Akpınar F. Entrapment of the sensory branch of the radial nerve (Wartenberg's syndrome): an unusual cause. *Tohoku J Exp Med.* 2001;193:251-4. [[Crossref](#)] [[PubMed](#)]
2. Marx SC, Dhalapathy S, Marx CA, et al. Ultrasonographical and histological cross-sectional study of the human superficial branch of the radial nerve. *Rom J Morphol Embryol.* 2011;52:1081-90. [[PubMed](#)]
3. Tzeng CY, Lee TS, Chen IC. Superficial radial nerve compression caused by a parosteal lipoma of proximal radius: a case report. *Hand Surg.* 2005;10:293-6. [[Crossref](#)] [[PubMed](#)]
4. Knutsen EJ, Calfee RP. Uncommon upper extremity compression neuropathies. *Hand Clin.* 2013;29:443-53. [[Crossref](#)] [[PubMed](#)]
5. Anthony JH, Hadeed A, Hoffler CE. Cheiralgia Paresthetica. 2021. In: *StatPearls [Internet]*. Treasure Island (FL): StatPearls Publishing; 2022. [[Link](#)]
6. Dang AC, Rodner CM. Unusual compression neuropathies of the forearm, part I: radial nerve. *J Hand Surg Am.* 2009;34:1906-14. [[Crossref](#)] [[PubMed](#)]
7. Braidwood AS. Superficial radial neuropathy. *J Bone Joint Surg Br.* 1975;57:380-3. [[Crossref](#)] [[PubMed](#)]
8. Meng S, Tinhofer I, Weninger WJ, et al. Anatomical and ultrasound correlation of the superficial branch of the radial nerve. *Muscle Nerve.* 2014;50:939-42. [[Crossref](#)] [[PubMed](#)]
9. Chodoroff G, Honet JC. Cheiralgia paresthetica and linear atrophy as a complication of local steroid injection. *Arch Phys Med Rehabil.* 1985;66:637-9. [[PubMed](#)]
10. Patel A, Pierce P, Chiu DTW. A fascial band implicated in Wartenberg syndrome. *Plast Reconstr Surg.* 2014;133:440e-2e. [[Crossref](#)] [[PubMed](#)]
11. Latef TJ, Bilal M, Vetter M, et al. Injury of the radial nerve in the arm: a review. *Cureus.* 2018;10:e2199. [[Crossref](#)] [[PubMed](#)] [[PMC](#)]
12. Kara M, Özçakar L, De Muyenck M, et al. Musculoskeletal ultrasound for peripheral nerve lesions. *Eur J Phys Rehabil Med.* 2012;48:665-74; quiz 708. [[PubMed](#)]
13. Visser LH. High-resolution sonography of the superficial radial nerve with two case reports. *Muscle Nerve.* 2009;39:392-5. [[Crossref](#)] [[PubMed](#)]